



## Large Bird Diet

Mazuri® Large Bird Diet is a nutritionally complete diet for large psittacines with diverse particle sizes to provide enrichment to the birds. This diet is specially formulated for non-breeding birds (young birds after first molt up to breeding age and birds kept in non-breeding situations). This diet is not for growing chicks.

### Features and Benefits

- **Nutritionally complete** – Requires no mixing or supplementation.
- **Extruded particle** means fewer fines and increased digestibility.
- **Better than seeds** – Less waste (20 to 50% of seed weight is hulls).
- **Reduced diet preparation** – Less time chopping fruits and veggies.
- **Formulation allows for supplementation with species-appropriate food items** – Can include up to 20% (by weight) of diet as produce, seeds, and other food items to encourage natural feeding behaviors and for training and enrichment.
- **Natural sources of pigments, antioxidants, and omega-3 fatty acids.**
- **No artificial colors or flavors.**

### Product Form & Packaging

Extruded Particle | 3 particle sizes

- **Catalog #0001455** | 25 lb. (11.33 kg) net weight paper w/poly bag
- **Catalog #0054031** | 3 lb. (1.36 kg) net weight resealable poly bag
- **Catalog #0053291** | Case: 6 x 3 lb. net weight resealable poly bags

### Guaranteed Analysis

Crude protein not less than .....	15.00%	Ash not more than .....	7.00%
Crude fat not less than.....	5.00%	Vitamin E not less than .....	150 IU/kg
Crude fiber not more than .....	6.00%	Omega-3 fatty acids not more than .....	0.60%
Moisture not more than .....	12.00%		

Total Microorganisms not less than .....270,000 CFU/g  
(Lactobacillus acidophilus, Lactobacillus casei, Bifidobacterium thermophilum, Enterococcus faecium)\*

\*Contains a source of live (viable) naturally occurring microorganisms.

### Ingredients

Ground Corn, Wheat Middlings, Dehulled Soybean Meal, Corn Gluten Meal, Ground Flaxseed, Soybean Oil, Dicalcium Phosphate, Calcium Carbonate, Wheat Germ, Glycerol Monostearate, Brewers Dried Yeast, Calcium Propionate (preservative), Salt, DL-Methionine, L-Lysine Dried Lactobacillus acidophilus Fermentation Product, L-Ascorbyl-2-Polyphosphate (Vitamin C), Dried Lactobacillus casei Fermentation Product, Choline Chloride, Dried Bifidobacterium thermophilum Fermentation Product, Biotin, Pyridoxine Hydrochloride, Dried Enterococcus faecium Fermentation Product, d-Alpha Tocopheryl Acetate (Vitamin E), Menadione Sodium Bisulfite Complex (Vitamin K), Vitamin D3 Supplement, Preserved with Mixed Tocopherols, Rosemary Extract, Folic Acid, Vitamin A Acetate, Citric Acid ( a Preservative), Canthaxanthin (color), Riboflavin Supplement, Zinc Oxide, Calcium Iodate, Manganous Oxide, Thiamine Mononitrate, Beta Carotene, Nicotinic Acid, Calcium Pantothenate, Vitamin B12 Supplement, Copper Sulfate, Zinc Sulfate, Sodium Selenite, Cobalt Carbonate.

### Feeding Directions

- Feed free choice to non-breeding birds and provide fresh food daily. A 1 lb. (450 g) bird will typically eat between 1/3 to 1/2 cup (33 – 40 g) per day, while a 2 lb. (900 g) bird may consume between 2/3 to 1 cup (66 – 80 g) per day.
- Fresh fruits, vegetables, and seeds may be provided, but at no more than 20% of the total diet.
- To switch birds from seeds (or their current diet) to this diet, changeover gradually. Start with 90% seed (or current diet) and 10% Mazuri® bird diet. On each succeeding day, reduce the amount of seeds by 10% while increasing the amount of the Mazuri® bird diet by 10%. On the 10th day, the diet should consist only of Mazuri® bird diet. Do not overfeed during this transition time.
  - Monitor the bird's weight and behavior carefully during the conversion period and make sure it is eating the new diet. Be especially attentive during the 10th to 12th day. If the bird is losing weight, or if the bird is not consuming the product, offer the bird a mix of 50% seed and 50% Mazuri® bird diet and restart the conversion process from this point.
  - If the bird is not eating, the excreta color will become dark green, and the volume will decrease. If the bird is consuming the bird diet, the excreta will be a brownish-green color.
- Always provide plenty of fresh, clean water. Thoroughly wash feed and water bowls on a regular basis. It is always good practice to wash hands thoroughly after feeding and/or handling animals. This diet is not for human consumption.

### Storage Conditions

For best results, reseal the bag between uses or store contents of open paper sack in container with sealing lid. Store in a cool (75°F/24°C or colder), dry (approximately 50% RH) location free from rodents and insects. Do not offer moldy or insect-infested feed to animals as it may result in illness, performance loss or death. Freezing will not harm the diet and may extend freshness. Use within 1 year of bag manufacturing or "Best if Used By" date.