



Emu Maintenance Diet

Mazuri® Emu Maintenance Diet is specifically formulated for emus during the non-breeding season and for yearlings selected as future breeders or replacement breeders.

Features and Benefits

- **High fiber and moderate energy diet** – Helps address obesity.
- **Balanced protein diet** – Helps to minimize amino acid imbalances and maximize protein use.
- **Contains vitamins known to be required by birds** – No supplementation necessary.
- **Naturally preserved with mixed tocopherols** – No ethoxyquin in formulation.

Product Form & Packaging

Catalog #0001488 | Pellet | 5/32" x 1/2" | 50 lb. (22.67 kg) paper w/poly bag

Guaranteed Analysis

Crude protein not less than	17.00%	Moisture not more than	12.00%
Crude fat not less than.....	2.00%	Ash not more than.....	9.00%
Crude fiber not more than.....	11.00%		

Ingredients

Ground Corn, Wheat Middlings, Dehydrated Alfalfa Meal, Dehulled Soybean Meal, Ground Soybean Hulls, Dicalcium Phosphate, Calcium Carbonate, Fish Meal, Cane Molasses, Salt, Calcium Lignin Sulfonate, DL-Methionine, Choline Chloride, Cholecalciferol, Pyridoxine Hydrochloride, DL-Alpha Tocopheryl Acetate (Form of Vitamin E), Biotin, Menadione Sodium Bisulfite Complex (source of Vitamin K), Vitamin A Acetate, Folic Acid, Preserved with Mixed Tocopherols (Form of Vitamin E), Calcium Pantothenate, Manganous Oxide, Rosemary Extract, Nicotinic Acid, Vitamin B-12 Supplement, Copper Sulfate, Riboflavin Supplement, Citric Acid, Thiamine Mononitrate, Zinc Oxide, Ferrous Carbonate, Zinc Sulfate, Calcium Iodate, Manganese Proteinate, Zinc Proteinate, Sodium Selenite, Cobalt Carbonate, Copper Proteinate.

Feeding Directions

- Feed free choice to emus selected for breeding.
 - From 6 – 8 months of age (about 70 – 90 lbs. or 31.8 – 40.8 kg body weight) until 2 weeks prior to the start of breeding season.
 - Two weeks prior to breeding season, transition birds to Mazuri® Ratite Breeder Diet.
 - Transition animals back to Mazuri® Emu Maintenance Diet towards the end of breeding season to accommodate for the large increase in feed consumption that occurs at this time.
- Always transition to new diets slowly – gradually transition over a 10-day period.
- Supplement high producing birds with calcium (using oyster shell) based on the level of egg production during the breeding season.
- Always provide plenty of fresh, clean water.
- Thoroughly wash feed and water bowls on a regular basis. It is always good practice to wash hands thoroughly after feeding and/or handling animals.
- This diet is not for human consumption.

Storage Conditions

For best results, reseal the bag between uses or store contents of open paper sack in container with sealing lid. Store in a cool (75°F/24°C or colder), dry (approximately 50% RH) location free from rodents and insects. Do not offer moldy or insect-infested feed to animals as it may result in illness, performance loss or death. Freezing will not harm the diet and may extend freshness. Use within 1 year of bag manufacturing or "Best if Used By" date.