



Ostrich Breeder

Mazuri® Ostrich Breeder Diet is a pelleted feed designed for breeding ostriches and rheas and is designed to be fed with a supplemental form of calcium, such as oyster shell.

Features and Benefits

- **Contains vitamins known to be required by birds** – Supplementation not required.
- **Contains high levels of added vitamin E and selenium** – Designed to address deficiencies in breeders and young chicks.
- **Naturally preserved with mixed tocopherols** – No ethoxyquin in formulation

Product Form & Packaging

Catalog #0001415 | Pellet | 5/32" x 1/2" | 50 lb. (22.67 kg) net weight paper w/poly bag

Guaranteed Analysis

Crude protein not less than.....	21.00%	Calcium not less than	1.65%
Crude fat not less than.....	2.50%	Calcium not more than	2.15%
Crude fiber not more than	11.00%	Phosphorus not less than.....	0.70%
Moisture not more than.....	12.00%	Sodium not more than	0.51%
Ash not more than	10.00%		

Ingredients

Dehulled Soybean Meal, Ground Corn, Dehydrated Alfalfa Meal, Ground Soybean Hulls, Wheat Middlings, Cane Molasses, Dicalcium Phosphate, Calcium Carbonate, Calcium Lignin Sulfonate, Fish Meal, Soybean Oil, Salt, Yeast Culture, Choline Chloride, DL-Methionine, Cholecalciferol, Biotin, DL-Alpha Tocopheryl Acetate (Form of Vitamin E), Pyridoxine Hydrochloride, Vitamin A Acetate, Menadione Sodium Bisulfite Complex (source of Vitamin K), Folic Acid, Thiamine Mononitrate, Preserved with Mixed Tocopherols (Form of Vitamin E), Calcium Pantothenate, Nicotinic Acid, Rosemary Extract, Vitamin B-12 Supplement, Riboflavin Supplement, Citric Acid, Copper Sulfate, Zinc Oxide, Manganous Oxide, Ferrous Carbonate, Manganese Proteinate, Zinc Sulfate, Zinc Proteinate, Calcium Iodate, Copper Proteinate, Sodium Selenite, Cobalt Carbonate.

Feeding Directions

- Ostrich: Feed free choice for very high-producing birds (over 100 eggs per year), at least 4 lbs. per bird per day during breeding season.
 - This feed may also be provided to birds that do not have access to pasture or have access to poor quality pasture.
 - Do not allow non-breeding birds to become overweight.
- Rhea & Cassowary: Feed free-choice from 3 months of age until 1 month before breeding.
- Always transition to new diets slowly (gradual transition over a 10-day period).
- Supplement high producing birds with calcium (using oyster shell). Mix the oyster shell with the feed based on the egg production during breeding season.

Egg production	Level of oyster shell to add (% of total diet)
from 0-30 eggs	No supplementation necessary
from 31-60 eggs	1%
from 61-90 eggs	2%
from 91-120 eggs	3%
from 121-150 eggs	4%

- Always provide plenty of fresh, clean water. Thoroughly wash feed and water bowls on a regular basis. It is always good practice to wash hands thoroughly after feeding and/or handling animals.
- This diet is not for human consumption.

Storage Conditions

For best results, reseal the bag between uses or store contents of open paper sack in container with sealing lid. Store in a cool (75°F/24°C or colder), dry (approximately 50% RH) location free from rodents and insects. Do not offer moldy or insect-infested feed to animals as it may result in illness, performance loss or death. Freezing will not harm the diet and may extend freshness. Use within 1 year of bag manufacturing or "Best if Used By" date.