



## Callitrichid NGA Gel Diet *(no glucose added)*

Mazuri® Callitrichid NGA Gel Diet is a palatable diet designed for diabetic marmosets, tamarins, and other callitrichids. This diet is shipped as a powder that is designed to be made into a gel prior to feeding.

### Features and Benefits

- **Designed with diabetic animals in mind** – Contains no added sucrose or glucose.
- **Convenient powder** – Mixes with hot water to form a palatable, soft, moist product.
- **Contains natural vitamin E.**
- **Meets NRC Non-Human Primate recommendations** – For vitamins, minerals, NDF, and ADF.

### Product Form & Packaging

Catalog #0047797 | Dry Powder | Available by the kg.

### Guaranteed Analysis (dry powder)

Crude protein not less than .....	20.00%	Ash not more than .....	7.00%
Crude fat not less than.....	7.50%	Selenium not less than .....	0.40 ppm
Crude fiber not more than .....	5.00%	Selenium not more than .....	0.48 ppm
Moisture not more than .....	10.00%		

### Ingredients

Corn Starch, Dehulled Soybean Meal, Maltodextrin, Ground Corn, Casein, Wheat Middlings, Gelatin, Corn Oil, Ground Wheat, Soybean Oil, Dried Egg Product, Fructose, Calcium Carbonate, Dried Whey, Dried Plain Beet Pulp, Corn Gluten Meal, Brewers Dried Yeast, Dehydrated Alfalfa Meal, Wheat Bran, Locust Bean Gum, Citric Acid, Natural Flavor, Carrageenan, Dicalcium Phosphate, Flaxseed Oil, Xanthan Gum, Dextrose, Salt, L-Ascorbyl-2-Polyphosphate (Vitamin C), DL-Methionine, Folic Acid, Choline Bitartrate, Maltodextrin, Aspartame, Taurine, Pyridoxine Hydrochloride, Cholecalciferol (Vitamin D3), Choline Chloride, Magnesium Oxide, L-Tryptophan, Zinc Proteinates, Iron Proteinates, Acesulfame Potassium, Ferrous Sulfate, D-Alpha Tocopheryl Acetate (Vitamin E), Zinc Sulfate, Manganese Proteinates, Calcium Pantothenate, Preserved with Mixed Tocopherols, Vitamin A Acetate, Menadione Sodium Bisulfite Complex (Vitamin K), Copper Proteinates, Potassium Carbonate, Citric Acid (a Preservative), Manganous Oxide, Thiamine Mononitrate, Zinc Oxide, Nicotinic Acid, Ferrous Carbonate, Vitamin B12 Supplement, L-Lysine, Riboflavin Supplement, Copper Sulfate, Cobalt Proteinates, Ethylenediamine Dihydroiodide, Rosemary Extract, Calcium Iodate, Sodium Selenite, Cobalt Carbonate, Biotin.

### Feeding Directions

NEVER FEED DRY PRODUCT.

- Combine by weight 60% boiling water and 40% Mazuri® gel powder. Adjust ratios to desired texture and need.
  - Mix fully, pour into shallow pan. Refrigerate until firm.
  - Cut to desired size and feed.
- Intake of up to 50% prepared Mazuri® gel is recommended.
  - Feed intake will vary based on age, body size and reproductive status.
  - A targeted intake level would be 2.5% - 3.0% powder per kg of animal by body weight per day (never feed dry powder without combining with water).
  - The gel diet can be fed in combination with fruit, vegetables, browse or other food items. Care should be taken when using supplemental dietary ingredients to ensure that nutrient recommendations are assessed.
- Always provide plenty of fresh, clean water.
- Prepared gel should be consumed or discarded within 24 hours or stored no longer than 7 days under refrigeration or up to 1 month frozen.
- If frozen, it is recommended that frozen batches be individually sealed in amount of usage, as repeated entry into the container exposes gel to oxygen.
- Thoroughly wash feed and water bowls on a regular basis. It is always good practice to wash hands thoroughly after feeding and/or handling animals.
- This diet is not for human consumption.

**Caution: Follow label directions: Feeding added selenium at levels in excess of 0.3 ppm in total diet is prohibited.**

### Storage Conditions

Mazuri® Gel Diets have a 1-year shelf life in the dry powder form when stored in a dry environment. For best results, tightly affix lid on canister after removal of desired dosage or store contents of open bags in container with sealing lid. Store in a cool (75°F/24°C or colder), dry (approximately 50% RH) location free from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death. The mixed product should be stored under refrigeration for no longer than 7 days or frozen for up to 1 month. It is recommended that frozen batches be individually sealed in amount of usage, as repeated entry into the container exposes gel to oxygen. Once this product is mixed, it should be handled like raw food.